IN THE UNITED STATES PATENT AND TRADEMARK OFFICE REQUEST FOR FILING NATIONAL PHARMACE TO STATE OF THE PROPERTY REQUEST FOR FILING NATIONAL PHAINTED TO 1 6 JUL 2001 PCT APPLICATION UNDER 35 U.S.C. 371 AND 37 CFR 1.494 OR 1.495

To:

Hon. Commissioner of Patents Washington, D.C. 20231



	MITTAL LETTER TO THE UNITED S		Atty Dkt:	P 0281578	/M99/0035/US				
DESIG	NATED/ELECTED OFFICE (DO/EO/	US)		M	# /Client Ref.				
From:	Pillsbury Winthrop LLP, IP Group:		Date: Ju	uly 16, 2001					
	This is a REQUEST for FILING a PO	CT/USA National I	Phase Applica	ition based on:					
1.	International Application	2. Internation	al Filing Date	3. Ea	arliest Priority Date Claimed				
	PCT/GB00/00237	28 Jan	uary 2000	5	February 1999				
•	û country code	Day M	ONTH Yea						
4.	Measured from the earliest priority difiled within:	late in item 3, this	PCT/USA Na		e item 2 if no earlier priority) oplication Request is being				
TOTAL	(a) 20 months from above item 3	date (b) 🖂 3	0 months fron	n above item 3	date,				
	(c) Therefore, the due date (unexter	ndable) isAugus	st 5, 2001	·					
	Title of Invention MEDICAMENT								
1	Inventor(s) BURNIE, James Pete	r et al							
Applica	nt herewith submits the following und	er 35 U.S.C. 371	to effect filing:						
7.1 8	☐ Please immediately start national	l examination prod	cedures (35 U	.S.C. 371 (f)).					
1965 1965 1965 1965 1965	☑ A copy of the International Application as filed (35 U.S.C. 371(c)(2)) is transmitted herewith (file if in English but, if in foreign language, file only if not transmitted to PTO by the International Bureau) including:								
	a. ☐ Request; b. ⋈ Abstract;								
	c. 24 pgs. Spec. and Claims; d sheet(s) Drawing which ar	re 🔲 informal 🔲	formal of size	☐ A4 ☐ 11	n n				
9.	☑ A copy of the International App	olication has bee	n transmitted	d by the Interna	ational Bureau.				
10.	A translation of the International								
	a. is transmitted herewith in (3) pgs. Spec. a	cluding: (1) R	equest; (2)] Abstract;					
		and Claims, awing which are:							
		informal [] form	nal of size] A4 □ 11"					
	b. is not required, as the ap								
	c. is not herewith, but will be Notice per Rule 494(c) if								
	d. Translation verification at			50x 7(b) 15 X U.					

RE: U	SA Natio	onal Phase Filing of PCT /GB00/00237
11.	\boxtimes	Please see the attached Preliminary Amendment Onal Phase Filing of PCT /GB00/00237 Please see the attached Preliminary Amendment JOTS Republic PCT/PTO 1 6 JUL 200
12.		Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)), i.e., <u>before 18th month</u> from first priority date above in item 3, are transmitted herewith (file only if in <u>English</u>) including:
13.	\boxtimes	PCT Article 19 claim amendments (if any) have been transmitted by the International Bureau
14.		Translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)), i.e., of claim amendments made before 18th month, is attached (required by 20th month from the date in item 3 if box 4(a) above is X'd, or 30th month if box 4(b) is X'd, or else amendments will be considered canceled).
15.	A dec a. ☐ b. ⊠	is submitted herewith
16.		ternational Search Report (ISR): s prepared by ⊠ European Patent Office □ Japanese Patent Office □ Other has been transmitted by the international Bureau to PTO. copy herewith (2 pg(s).) ☑ plus Annex of family members (1 pg(s).).
	Intern a.	during Examination) including attached amended:
18.	Inform a. 🛭 b. 🖾 c. 🖾	Attached Form PTO-1449 listing documents Attached copies of documents listed on Form PTO-1449 A concise explanation of relevance of ISR references is given in the ISR. Assignment document and Cover Sheet for recording are attached. Please mail the recorded assignment document back to the person whose signature, name and address appear at the end of this letter.
20.		Copy of Power to IA agent.
21.		Drawings (complete only if 8d or 10a(4) not completed): sheet(s) per set: ☐ 1 set informal; ☐ Formal of size ☐ A4 ☐ 11"
22. 22(a)		Entity Status is Not claimed is claimed (pre-filing confirmation required) (No.) Small Entity Statement(s) enclosed (since 9/8/00 Small Entity Statements(s) not essential to claim)
(1) (3) (5)	filed in in (cou	(4) (6) See Form PCT/IB/304 sent to US/DO with copy of priority documents. If copy has not been received, please proceed promptly to obtain same from the IB.

24. Attached: 9 pages of Sequence Listiing

25	Per	Item	17.c2, cancel origi	nal page	es #, clai	ms #, Drav	ving Sheets	#	-		
26. Based	Calc on an	culat nend	ion of the U.S. Nat ed claim(s) per abo	ional Fe	e (35 U.S.C. 37	71 (c)(1)) and ot 4,	her fees is a (hilite)	s follo	ws:		
Total E Indeper	ndent	Clai		ole Depe	minus 20 = minus 3 = ndent claim is į	present,	x \$18/\$9 x \$80/\$40 add\$270/\$1	=	\$0 \$0 +0	964	/967 /965 /969
BASIC	NATI	ONA	L FEE (37 CFR 1.4	92(a)(1)-	(4)): →→ BAS	SIC FEE REQUIR	ED, NOW -	> >>	т →		
A.	If co	untry	code letters in item	1 are <u>n</u>	ot "US","BR","E	BB","TT","MX","IL	" "NZ", "IN" c	or "ZA"	. ↓ Ji		
	See 1. 2.	Sea	<u>16 re:</u> Irch Report was <u>not</u> Irch Report was pre	prepared pared by	d by EPO or JF EPO or JPO -	<u> </u>	add\$1000/\$ add\$860/\$4	5500 130 _	+430		60/961 970/971
SKIP B,	C, D	AND	E UNLESS country	code lette	ers in item 1 are	e "US","BR","BB"	,"TT","MX","	IL", "N	Z", "IN"	or "ZA"	
**************************************		B.	If <u>USPTO</u> did not i (ISR) <u>and</u> (if box 4 Examination Repo	ssue <u>bot</u> (b) abov	h International e is X'd) the Int	Search Report ternational	add\$1000/\$		+0		60/961
(<u>ontv)</u> (<u>one)</u> → (<u>off)</u> (these)		C.	If <u>USPTO</u> issued I	SR but n	ot IPER (or bo	x 4(a) above is	add\$710/\$3	355 -	+0	9	58/959
(<u>+</u> 4) → (<u>boxes</u>)		D.	If <u>USPTO</u> issued I YES,	PER but	IPER Sec. V b	ooxes <u>not all</u> 3	add\$690/\$3	³⁴⁵ -	+0	9	56/957
		E.	If international pre USPTO and Rules Sec. V all 3 boxes	s 492(a)(4) and 496(b) s	satisfied (IPER	add \$100/\$	50 <u>-</u>	+0	9	62/963
							SUBTOTAI	L = =	\$430		
28.		Ū	ment box 19 above	ŕ	· ·	_		_	+0	(5	581)
29.	Atta	ched	is a check to cover	the			TOTAL FE	ES =	\$430		
			osit Account No. 03- er No. 05088		0281578 M#			090			
filed, or who or hereafted duplicate of	ich shou r relative opy of tl	uld have e to this his shee	The Commissioner is hereby be been filed herewith or conce application and the resulting et is attached. ENT does not authorize char	ming any par Official docur	er filed hereafter, and nent under Rule 20, or	which may be required ur r credit any overpayment,	nder Rules 16-18 a to our Account/Ord	nd 492 (<u>m</u>	issing or ins	ufficient fee on	l <u>y</u>) now
THIS OTHER	(OL OI	AT LINE	<u>uoes noi authorize</u> ondi	Pillsbu	ry Winthrop L tual Property	.LP					
				By Atty	Paul N. Ko	okulis		Reg.	No′	16773	
Atty/Se	c: PN	K/mh	nn ,	Sig:	De	MIL	7	Fax: Tel:		703) 905-2 703) 905-2	
	N	JOTE	File in dunlicate w	ith 2 nos	toard receipts	(PAT-103) & atta	chments				

Rec'd PCT/PTO 20 NOV 2001

Attorney Docket No.

Applicant or Patentee:

James Peter BURNIE and Ruth Christine MATTHEWS

International Appl. No.: Filed:

PCT/GB00/00237 28 January 2000

For:

Medicament

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS (37 CFR 1.9(f) and 1.27(b)) INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 CFR 1.9 (c) for purposes of paying reduced fees under section 41 (a) and (b) of Title 35, United States Code, to the Patent and Trademark Office with regard to the invention entitled described in:

[] the specification filed herewith

[X] International application no. PCT/GB00/00237

[X] filed 28 January 2000

I have not assigned, granted, conveyed or licensed except as shown in the attachment hereto and am under no obligation under contract or law to assign, grant, convey or license, any rights in the invention to any person who could not be classified as an independent inventor under 37 CFR 1.9 (c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 CFR 1.9 (d) or a nonprofit organization under 37 CFR 1.9 (e).

Each person, concern or organization to which I have assigned, granted, conveyed, or licensed or am under an obligation under contract or law to assign, grant, convey, or license any rights in the invention is listed below:

[] no such person, concern, or organization

[X] persons, concerns or organizations listed

below*

FULL NAME: ADDRESS:

NeuTec Pharma plc St James's Court

Brown Street Manchester

M2 2JT Great Britain

T 1

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28 (b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

[]

Date

Ruth Christine MATTHEWS

11 July 01

Rec'd PCT/PTO 2 0 NOV 2001 Attorney Docket No.

Applicant or Patentee James Peter BURNIE and Ruth Christine MATTHEWS

Serial or Patent No.: PCT/GB00/00237
Filed or Issued: 20 January 2000
Title: Medicament

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS [37 CFR 1.9 (f) AND 1.27 (c)] - SMALL BUSINESS CONCERN

1 nereby	declare that I a	m		

[X] the owner of the small business concern identified below:[] an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF CONCERN NeuTec Pharma plc

ADDRESS OF CONCERN St James's Court

Brown Street Manchester M2 2JT GB

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 CFR 121.3-18 and reproduced in 37 CFR 1.9(d), for purpose of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties control or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention, entitled

Medicament

i in

[]	the specification file	d herewith	
[X]	the application ident	ified above	
[]	Application Serial N	0.	filed
[]	Patent No.	, issued	

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under

statements are required from each named person, concern, or organization having rights to the invention averring to their status as small entities [37 CFR 1.27].
FULL NAME ADDRESS [] INDIVIDUAL [] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION
FULL NAME ADDRESS [] INDIVIDUAL [] SMALL BUSINESS CONCERN [] NONPROFIT ORGANIZATION
I acknowledge the duty to file, in this Application or Patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate [37 CFR 1.28(b)].
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.
NAME OF PERSON SIGNING James Sume
TITLE OF PERSON OTHER THAN OWNER DURCLET
ADDRESS OF PERSON SIGNING 1 Greystate Drive Chesleme
SIGNATURE DATE! July 1

37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e). *Note: Separate verified

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re PATENT APPLICATION OF

Inventor(s): BURNIE, James Peter et al

Filed: Herewith

Title: MEDICAMENT

July 16, 2001

PRELIMINARY AMENDMENT

Hon. Commissioner of Patents Washington, D.C. 20231 Sir: Please amend this application as follows: IN THE SPECIFICATION: At the top of the first page, just under the title, insert -- This application is the National Phase of International Application \boxtimes PCT/GB00/00237 filed January 28, 2000 which designated the U.S. and that International Application was not published under PCT Article 21(2) in English.-was was Respectfully submitted, PILLSBURY WINTHROP LLP Intellectual Property Group

Attorney: Paul N. Kokulis

Reg. No: 16773

Tel. No.: (703) 905-2118 Fax No.: (705) 905-2500

Atty\Sec. PNK/mhn 1600 Tysons Boulevard

McLean, VA 22102 (703) 905-2000

PCT/GB00/00237

-1-

JC18 Rec'd PCT/PTO 1 6 JUL 2001

SEQUENCE LISTING

<110> NeuTec Pharma plc	
<120> Medicament	
<130> M99/0035/PCT	
<140><141>	
<150> GB9902555.3 <151> 1999-02-05	
<160> 16	
<170> PatentIn Ver. 2.1	
<210> 1 <211> 1491 <212> DNA <213> Chlamydia pneumoniae	
<220> <221> CDS	
<222> (1)(1491)	
<pre><400> 1 gat aca aac atg tct att tca tct tct tca gga cct gac aat caa aaa Asp Thr Asn Met Ser Ile Ser Ser Ser Gly Pro Asp Asn Gln Lys</pre>	48
aat atc atg tct caa gtt ctg aca tcg aca ccc cag ggc gtg ccc caa Asn Ile Met Ser Gln Val Leu Thr Ser Thr Pro Gln Gly Val Pro Gln 20 25 30	96
caa gat aag ctg tct ggc aac gaa acg aag caa ata cag caa aca cgt Gln Asp Lys Leu Ser Gly Asn Glu Thr Lys Gln Ile Gln Gln Thr Arg 35 40 45	144
cag ggt aaa aac act gag atg gaa agc gat gcc act att gct ggt gct Gln Gly Lys Asn Thr Glu Met Glu Ser Asp Ala Thr Ile Ala Gly Ala 50 55 60	192
tct gga aaa gac aaa act tcc tcg act aca aaa aca gaa aca gct cca Ser Gly Lys Asp Lys Thr Ser Ser Thr Thr Lys Thr Glu Thr Ala Pro 65 70 75 80	240
caa cag gga gtt gct gct ggg aaa gaa tcc tca gaa agt caa aag gca Gln Gln Gly Val Ala Ala Gly Lys Glu Ser Ser Glu Ser Gln Lys Ala 85 90 95	288
ggt gct gat act gga gta toa gga gcg gct gct act aca gca toa aat	336

Gly	Ala	Asp	Thr 100	Gly	Val	Ser	Gly	Al a 105	Ala	Ala	Thr	Thr	Ala 110	Ser	Asn	
										att Ile					aaa Lys	384
										agc Ser					caa Gln	432
atg Met 145	aaa Lys	gaa Glu	gtc Val	gaa Glu	gcg Ala 150	gtt Val	gtt Val	gtt Val	gct Ala	gcc Ala 155	ctc Leu	tca Ser	gly aaa	aaa Lys	agt Ser 160	480
										ctc Leu						528
aca Thr	cca Pro	aga Arg	tca Ser 180	gag Glu	gtt Val	atc Ile	gaa Glu	atc Ile 185	gga Gly	ctc Leu	gcg Ala	ctt Leu	gct Ala 190	aaa Lys	gca Ala	576
										gcc Ala						624
										aaa Lys					aag Lys	672
										gaa Glu 235						720
										gga Gly					gtc Val	768
										att Ile					att Ile	816
gtt Val	gct Ala	gct Ala 275	att Ile	ttt Phe	aca Thr	tgc Cys	gga Gly 280	gct Ala	gga Gly	ctc Leu	gct Ala	gga Gly 285	cto Leu	gct Ala	gcg Ala	864
gga Gly	gct Ala 290	gct Ala	gta Val	ggt Gly	gca Ala	gcg Ala 295	gca Ala	gct Ala	gga Gly	ggt Gly	gca Ala 300	gca Ala	gga Gly	gct Ala	gct Ala	912
										gtt Val					caa Gln	960

WO 00/46359 PCT/GB00/00237

- 3 -

305	310	315	320
gcg gtg aaa caa gct Ala Val Lys Gln Ala 325	gtt atc aca gct gtc Val Ile Thr Ala Val 330	aga caa gcg atc ac Arg Gln Ala Ile Th	r Ala
gct ata aaa gcg gct Ala Ile Lys Ala Ala 340	gtc aaa tct gga ata Val Lys Ser Gly Ile 345	aaa gca ttt atc aa Lys Ala Phe Ile Lys 350	a act 1056 s Thr
tta gtc aaa gcg att Leu Val Lys Ala Ile 355	gcc aaa gcc att tct Ala Lys Ala Ile Ser 360	aaa gga atc tct aa Lys Gly Ile Ser Lys 365	g gtt 1104 s Val
ttc gct aag gga act Phe Ala Lys Gly Thr 370	caa atg att gcg aag Gln Met Ile Ala Lys 375	aac ttc ccc aag ct Asn Phe Pro Lys Let 380	c tcg 1152 1 Ser
aaa gtc atc tcg tct Lys Val Ile Ser Ser 385	ctt acc agt aaa tgg Leu Thr Ser Lys Trp 390	gtc acg gtt ggg gt Val Thr Val Gly Val 395	t ggg 1200 l Gly 400
gtt gta gtt gcg gcg Val Val Val Ala Ala 405	cct gct ctc ggt aaa Pro Ala Leu Gly Lys 410	ggg att atg caa at Gly Ile Met Gln Met 41!	Gln
ctc tcg gag atg caa Leu Ser Glu Met Gln 420	caa aac gtc gct caa Gln Asn Val Ala Gln 425	ttt cag aaa gaa gt Phe Gln Lys Glu Val 430	c gga 1296 L Gly
aaa ctg cag gct gcg Lys Leu Gln Ala Ala 435	gct gat atg att tct Ala Asp Met Ile Ser 440	atg ttc act caa tt Met Phe Thr Gln Phe 445	t tgg 1344 E Trp
caa cag gca agt aaa Gln Gln Ala Ser Lys 450	att gcc tca aaa caa Ile Ala Ser Lys Gln 455	aca ggc gag tct aa Thr Gly Glu Ser Asr 460	t gaa 1392 1 Glu
atg act caa aaa gct Met Thr Gln Lys Ala 465	acc aag ctg ggc gct Thr Lys Leu Gly Ala 470	caa atc ctt aaa gc Gln Ile Leu Lys Ala 475	g tat 1440 a Tyr 480
gcc gca atc agc gga Ala Ala Ile Ser Gly 485	gcc atc gct ggc gca Ala Ile Ala Gly Ala 490	cat aaa acc aat aa His Lys Thr Asn Ass 495	n Phe
taa			1491

<210> 2

<211> 496

<212> PRT

<213> Chlamydia pneumoniae

<400> 2 Asp Thr Asn Met Ser Ile Ser Ser Ser Ser Gly Pro Asp Asn Gln Lys 10 Asn Ile Met Ser Gln Val Leu Thr Ser Thr Pro Gln Gly Val Pro Gln Gln Asp Lys Leu Ser Gly Asn Glu Thr Lys Gln Ile Gln Gln Thr Arg 40 Gln Gly Lys Asn Thr Glu Met Glu Ser Asp Ala Thr Ile Ala Gly Ala 55 60 Ser Gly Lys Asp Lys Thr Ser Ser Thr Thr Lys Thr Glu Thr Ala Pro 70 Gln Gln Gly Val Ala Ala Gly Lys Glu Ser Ser Glu Ser Gln Lys Ala Gly Ala Asp Thr Gly Val Ser Gly Ala Ala Ala Thr Thr Ala Ser Asn 105 Thr Ala Thr Lys Ile Ala Met Gln Thr Ser Ile Glu Glu Ala Ser Lys 1115 120 Ser Met Glu Ser Thr Leu Glu Ser Leu Gln Ser Leu Ser Ala Ala Gln 135 140 Met Lys Glu Val Glu Ala Val Val Ala Ala Leu Ser Gly Lys Ser 150 155 Ser Gly Ser Ala Lys Leu Glu Thr Pro Glu Leu Pro Lys Pro Gly Val 165 170 Thr Pro Arg Ser Glu Val Ile Glu Ile Gly Leu Ala Leu Ala Lys Ala 185 Ile Gln Thr Leu Gly Glu Ala Thr Lys Ser Ala Leu Ser Asn Tyr Ala 200 Ser Thr Gln Ala Gln Ala Asp Gln Thr Asn Lys Leu Gly Leu Glu Lys 215 220 Gln Ala Ile Lys Ile Asp Lys Glu Arg Glu Glu Tyr Gln Glu Met Lys 235 230 Ala Ala Glu Gln Lys Ser Lys Asp Leu Glu Gly Thr Met Asp Thr Val 245 250 Asn Thr Val Met Ile Ala Val Ser Val Ala Ile Thr Val Ile Ser Ile 265 Val Ala Ala Ile Phe Thr Cys Gly Ala Gly Leu Ala Gly Leu Ala Ala Gly Ala Ala Val Gly Ala Ala Ala Gly Gly Ala Ala Gly Ala Ala 295 300 Ala Ala Thr Thr Val Ala Thr Gln Ile Thr Val Gln Ala Val Val Gln 315 310 Ala Val Lys Gln Ala Val Ile Thr Ala Val Arg Gln Ala Ile Thr Ala 325 330 Ala Ile Lys Ala Ala Val Lys Ser Gly Ile Lys Ala Phe Ile Lys Thr 345 Leu Val Lys Ala Ile Ala Lys Ala Ile Ser Lys Gly Ile Ser Lys Val 360 Phe Ala Lys Gly Thr Gln Met Ile Ala Lys Asn Phe Pro Lys Leu Ser 375 380 Lys Val Ile Ser Ser Leu Thr Ser Lys Trp Val Thr Val Gly Val Gly 390 395

 Val
 Val
 Ala
 Ala
 Pro
 Ala
 Leu
 Gly
 Lys
 Gly
 Ile
 Met
 Gln
 Met
 Gln
 Asn
 Val
 Ala
 Gln
 Phe
 Gln
 Lys
 Glu
 Val
 Gly
 Asn
 Val
 Ala
 Gln
 Phe
 Gln
 Lys
 Glu
 Val
 Gly
 Asn
 Asn
 Ala
 Ala
 Ala
 Asn
 Met
 Ile
 Ser
 Met
 Ile
 Ser
 Met
 Ile
 Asn
 Frage
 Asn
 Gly
 Ala
 Asn
 Gly
 Ala
 Asn
 Gly
 Ala
 Asn
 Gly
 Asn
 <td

<210> 3

<211> 302 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Codon optimised N-terminal section of Chlamydia pneumoniae protein

<220>

<221> UNSURE

<222> (1)..(30)

<223> S-tag and thrombin cleavage site

<220>

<221> UNSURE

<222> (292)..(302)

<223> Histidine tag

<400> 3

Met Lys Glu Thr Ala Ala Ala Lys Phe Glu Arg Gln His Met Asp Ser 1 5 10 15

Pro Asp Leu Gly Thr Leu Val Pro Arg Gly Ser Ala Ile Ser Asp Pro 20 25 30

Asp Thr Asn Met Ser Ile Ser Ser Ser Ser Gly Pro Asp Asn Gln Lys
35 40 45

Asn Ile Met Ser Gln Val Leu Thr Ser Thr Pro Gln Gly Val Pro Gln 50 55 60

Gln Asp Lys Leu Ser Gly Asn Glu Thr Lys Gln Ile Gln Gln Thr Arg
65 70 75 80

Gln Gly Lys Asn Thr Glu Met Glu Ser Asp Ala Thr Ile Ala Gly Ala 85 90 95 Ser Gly Lys Asp Lys Thr Ser Ser Thr Thr Lys Thr Glu Thr Ala Pro 100 105 110

Gln Gln Gly Val Ala Ala Gly Lys Glu Ser Ser Glu Ser Gln Lys Ala 115 120 125

Gly Ala Asp Thr Gly Val Ser Gly Ala Ala Ala Thr Thr Ala Ser Asn 130 135 140

Thr Ala Thr Lys Ile Ala Met Gln Thr Ser Ile Glu Glu Ala Ser Lys 145 150 155 160

Ser Met Glu Ser Thr Leu Glu Ser Leu Gln Ser Leu Ser Ala Ala Gln 165 170 175

Met Lys Glu Val Glu Ala Val Val Ala Ala Leu Ser Gly Lys Ser 180 185 190

Ser Gly Ser Ala Lys Leu Glu Thr Pro Glu Leu Pro Lys Pro Gly Val
195 200 205

Thr Pro Arg Ser Glu Val Ile Glu Ile Gly Leu Ala Leu Ala Lys Ala 210 215 220

Ile Gln Thr Leu Gly Glu Ala Thr Lys Ser Ala Leu Ser Asn Tyr Ala
225 230 235 240

Ser Thr Gln Ala Gln Ala Asp Gln Thr Asn Lys Leu Gly Leu Glu Lys 245 250 255

Gln Ala Ile Lys Ile Asp Lys Glu Arg Glu Glu Tyr Gln Glu Met Lys 260 265 270

Ala Ala Glu Gln Lys Ser Lys Asp Leu Glu Gly Thr Met Asp Thr Val 275 280 285

Asn Thr Val Ala Ala Ala Leu Glu His His His His His His 290 295 300

<210> 4

<211> 9

<212> PRT

<213> Chlamydia pneumoniae

<400> 4

Ser Ala Lys Leu Glu Thr Pro Glu Leu

<210> 5

<211> 7

<212> PRT

```
<213> Chlamydia pneumoniae
<400> 5
Pro Lys Pro Gly Val Thr Pro
<210> 6
<211> 9
<212> PRT
<213> Chlamydia pneumoniae
Gly Val Thr Pro Arg Ser Glu Val Ile
<210> 7
<211> 6
<212> PRT
<213> Chlamydia pneumoniae
<400> 7
Glu Val Ile Glu Ile Gly
<210> 8
<211> 8
<212> PRT
<213> Chlamydia pneumoniae
<400> 8
Ala Ile Lys Ile Asp Lys Glu Arg
<210> 9
<211> 6
<212> PRT
<213> Chlamydia pneumoniae
<400> 9
Ser Gly Pro Asp Asn Gln
<210> 10
<211> 9
<212> PRT
<213> Chlamydia pneumoniae
<400> 10
```

```
Ser Gly Asn Glu Thr Lys Gln Ile Gln
 1
                  5
<210> 11
<211> 9
<212> PRT
<213> Chlamydia pneumoniae
<400> 11
Ser Glu Gly Gln Lys Ala Gly Ala Asp
<210> 12
<211> 9
<212> PRT
<213> Chlamydia pneumoniae
<400> 12
Thr Ala Ile Glu Glu Ala Ser Lys Ser
<210> 13
<211> 9
<212> PRT
<213> Chlamydia pneumoniae
<400> 13
Ser Lys Ser Met Glu Ser Thr Leu Glu
<210> 14
<211> 9
<212> PRT
<213> Chlamydia pneumoniae
 <400> 14
 Glu Tyr Gln Glu Met Lys Ala Ala Glu
 <210> 15
 <211> 14
 <212> PRT
 <213> Chlamydia pneumoniae
 <400> 15
 Glu Lys Gln Ala Ile Lys Ile Asp Lys Glu Arg Glu Glu Tyr
                   5
```

<210> 16
<211> 14
<212> PRT
<213> Chlamydia pneumoniae

<400> 16
Glu Thr Pro Glu Leu Pro Lys Pro Gly Val Thr Pro Arg Ser
1 5 10

APPLICATION UNDER UNITED STATES PATENT LAWS 200

Atty. Dkt. No. PW 0281578/M99/0035/US

(M#)

Invention:

MEDICAMENT

Inventor (s):

BURNIE, James Peter

MATTHEWS, Ruth Christine

Pillsbury Winthrop LLP Intellectual Property Group 1600 Tysons Boulevard

McLean, VA 22102 Attorneys

This is a:

Sub. Spec Filed

Sub. Spec. filed

in App. No.

Marked up Specification re

In App. No

Telephone: (703) 905-2000

	Provisional Application
	Regular Utility Application
	Continuing Application ☑ The contents of the parent are incorporated by reference
\boxtimes	PCT National Phase Application
	Design Application
	Reissue Application
	Plant Application
	Substitute Specification

SPECIFICATION

PCT/GB00/00237

<u>Medicament</u>

The present invention concerns treatment, prevention and diagnosis of infection due to *Chlamydia pneumoniae* and in particular to the prevention and treatment of atherosclerosis, including coronary atherosclerosis, caused by same.

C. pneumoniae is associated with atherosclerosis but no definitive link between the two has yet been established (Hammerschlag, M.R., 1998, Eur. J. Clin. Microbiol. Infect. Dis., 17: 305-308). Friedank, H.M. et al. (1993, Eur. J. Clin. Microbiol. Infect. Dis., 12(12): 947-951) identify a 54 kDa C. pneumoniae antigen which was recognised by 93% of sera positive for C. pneumoniae, the antigen appearing to be located on the surface of elementary bodies. Wiedman, A.A.M. et al. (1997, Clin. Diagn. Labs. Immunol., 4(6):700-704) showed the infectivity of C. pneumoniae elementary bodies to be slightly reduced by the use of antibody specific against a 54 kDa C. pneumoniae protein.

Despite investigating it, other researchers have not confirmed the immunogenicity of the *C. pneumoniae* 54 kDa band (see for example Kutlin, A. and Roblin, P.M., 1998, J. Infect. Dis., 177: 720-724; Campbell, L.A. et al., 1990, J. Clin. Microbiol., 28(6): 1261-1264; Campbell, L.A. et al., 1990. Infection and Immunity, 58(1): 93-97; Puolakkainen, M. et al., 1993, J. Clin. Microbiol., 31(8): 2212-2214; hkima, Y. et al., 1994, J. Clin. Microbiol., 32(3): 583-588; Maass, M. and Gieffers, J., 1997, J. Infection, 35: 171-176; Gonen, R. et al., 1993, APMIS, 101:719-726).

The present inventor has now succeeded in isolating, purifying and identifying a C. pneumoniae protein which (together with inhibitors of same, such as

WO 00/46359 PCT/GB00/00237

-2-

antibodies) is protective and therapeutic against *C. pneumoniae* infection. The therapeutic role of the protein has previously neither been suggested nor disclosed.

According to the present invention there is provided a *C. pneumoniae* protein having the amino acid sequence of SEQ ID NO: 2, for use in a method of treatment or diagnosis of the human or animal body. The amino acid sequence has been confirmed by N-terminal amino-acid sequencing (see "Experimental" below) and the protein has a theoretical molecular weight of 50.8 kDa, although post-translational modifications such as glycosylation may of course affect its apparent molecular weight as determined by e.g. SDS-PAGE. Experiments (below) have shown it to have an apparent molecular weight of 51 kDa on SDS-PAGE gels.

As can be seen from the plethora of publications above, although some identify immunogenic bands at molecular weights of 50-54 kDa, no specific therapeutically effective proteins have been identified.

Experiments (below) have allowed the present inventor to isolate and purify the protein of the present invention and identify the gene sequence coding for the protein. This has allowed the determination of the protein amino acid sequence (above). The nucleotide sequence coding for same forms another part of the present invention. Thus according to the present invention there is also provided a nucleotide sequence coding for a protein according to the present invention, for use in a method of treatment or diagnosis of the human or animal body. Such a nucleotide sequence may have the sequence of SEQ ID NO: 1. Modified nucleotide sequences having codons encoding the same amino acid sequence will be readily apparent to one skilled in the art.

The nucleotide sequence of the present invention and the amino acid sequence it encodes are already known from the Chlamydia Genome Project

WO 00/46359 PCT/GB00/00237

- 3 -

(C. pneumoniae CWL029/CPn0809), as is an apparent C. trachomatis homologue (CT578). However, therapeutic and diagnostic uses for same have not been previously suggested.

The invention also extends to encompass forms of the protein which have been insubstantially modified (i.e. which have been partially modified), particularly forms of the protein which display the same immunogenic properties as the protein itself.

By "partial modification" and "partially modified" is meant, with reference to amino acid sequences, a partially modified form of the molecule which retains substantially the properties of the molecule from which it is derived, although it may of course have additional functionality. Partial modification may, for example, be by way of addition, deletion or substitution of amino acid residues. Substitutions may be conserved substitutions. Hence the partially modified molecule may be a homologue of the molecules from which it was derived. It may, for example, have at least 70% homology with the molecule from which it was derived. It may for example have at least 80, 90 or 95% homology with the molecule from which it was derived. An example of a homologue is an allelic mutant.

Also provided according to the present invention is the use of a protein, immunogenic fragment thereof or nucleic acid sequence encoding same according to the present invention in the manufacture of a medicament for the treatment of infection due to *C. pneumoniae*.

Immunogenic fragments of the protein include any fragment of the protein which elicits an immune response, and includes epitopes. Analogues (mimotopes) of epitopes may be readily created, the mimotopes having different sequences but displaying the same epitope and thus the term "immunogenic fragments" also

encompasses immunogenic analogues of the fragments e.g. mimotopes. Epitopes may be readily determined and mimotopes readily designed (Geysen, H.M. et al., 1987, Journal of Immunological Methods, 102: 259-274; Geysen, H.M. et al., 1988, J. Mol. Recognit., 1(1):32-41; Jung, G. and Beck-Sickinger, A.G., 1992, Angew. Chem. Int. Ed. Eng., 31: 367-486). Such an immunogenic fragment carrying epitopes may also be described as being a peptide having the amino acid sequence of the immunogenic fragment and which carries an epitope.

The present inventor has succeeded in isolating a number of epitopes (immunogenic fragments) of the protein of the present invention. Thus according to the present invention there is also provided an epitope having the amino acid sequence of any one of SEQ ID NOs: 4-14. In particular, SEQ ID NOs: 5-7 provide an overlapping set of highly immunogenic peptides - as can be seen from the experimental data (below) SEQ ID NO: 5 provides for especially good results. Similarly, excellent results are also obtained from SEQ ID NO: 8.

The protein, immunogenic fragments thereof and nucleic acid sequences encoding same may be used in therapy, both prophylactically (e.g. as immunostimulants such as vaccines) and for treatment of infection due to *C. pneumoniae*. For example a nucleotide sequence encoding the protein or immunogenic fragment thereof may be used in the manufacture of a DNA vaccine (Montgomery, D.L. *et al.*, 1997, Pharmacol. Ther., 74(2): 195-205; Donnelly, J.J. *et al.*, 1997, Annu. Rev. Immunol., 15: 617-648; Manickan, E. *et al.*, 1997, Crit. Rev. Immunol., 17(2): 139-154).

Binding agents and inhibitors (such as antibodies or other neutralising agents) specific against the protein and immunogenic fragments thereof may also be used both diagnostically and therapeutically. Binding agents have a target to which they are specific, and in the case of a binding agent being an antibody, the target is an antigen.

WO 00/46359 PCT/GB00/00237

- 5 -

An example of a therapeutic medicament is antibody specific against the protein of the present invention, and this may be employed in immunotherapy, for example passive immunotherapy. Antibodies, their manufacture and use are well known (Harlow, E. and Lane, D., "Using Antibodies - A Laboratory Manual", Cold Spring Harbor Laboratory Press, New York, 1998) and so antibodies and antigen binding fragments thereof will be readily apparent to one skilled in the art, and reference herein to antibodies is also reference to antigen binding fragments unless stated otherwise. Other inhibitors such as ribozymes, antisense oligonucleotides and DNA vaccines will be readily apparent to one skilled in the art (Fries, P.C., 1999, "DNA Vaccines", New England Journal of medicine, 341: 1623-1624; Leitner, W.W. et al., 1999, "DNA and RNA based vaccines: principles, progress and prospects", Vaccine, 18: 765-777; Muotri, A.R. et al., 1999, "Ribozymes and the anti-gene therapy: how a catalytic RNA can be used to inhibit gene function", Gene, 237: 303-310; Rossi, J.J., 1999, "Ribozymes, genomics and therapeutics", Chemistry & Biology, 6: R33-R37; James, H.A., 1999, "The potential application of ribozymes for the treatment of haematological disorders", Journal of Leukocyte Biolofy, <u>66</u>: 361-368)

Thus the present invention also provides the use of a inhibitor specific to the protein of the present invention in the manufacture of a medicament for the treatment of infection due to *C.pneumoniae*.

Also provided according to the present invention is a method of manufacture of a medicament for the treatment of infection due to *C. pneumoniae*, characterised in the use of a protein, immunogenic fragment or inhibitor according to the present invention.

Also provided according to the present invention is a method of treatment of infection due to *C. pneumoniae*(e.g. of a patient in need of same), comprising the step

of administering to a patient a medicament comprising a protein, immunogenic fragment or inhibitor according to the present invention. The exact dose of medicament administered to a patient may be readily determined using simple dose-response assays. Medicaments may additionally comprise a pharmaceutically acceptable carrier, diluent or excipient (Remington's Pharmaceutical Sciences and US Pharmacopeia, 1984, Mack Publishing Company, Easton, PA, USA)

It has not been previously suggested that the protein of the present invention (or immunogenic fragments of same) is diagnostic for infection due to *C. pneumonia*. Binding agents specific to the protein of the present invention (for example antibodies) may also be used diagnostically, for example in an ELISA-type test. Thus also provided according to the present invention is the use of a protein, immunogenic fragment or binding agent according to the present invention in the manufacture of a diagnostic test for *C. pneumoniae*.

Also provided is a diagnostic test method for infection due to C. pneumoniae comprising the steps of:

- I) reacting an antibody specific against the protein of the present invention with serum from a patient;
 - ii) detecting an antibody-antigen binding reaction; and
- iii) correlating the detection of an antibody-antigen binding reaction with the presence of the protein.

Such test methods may also be performed using other binding agents specific to the protein of the present invention.

Also provided is a kit of parts for performing such a test, characterised in that it comprises antibody specific against the protein of the present invention.

The invention will be further apparent from the following description, with reference to the several figures of the accompanying drawings, which show, by way of example only, uses of the proteins of the present invention.

EXPERIMENTAL

The experiments below detail the identification of a number of peptides and antisera against same which are useful in the therapy and diagnosis of infections due to *Chlamydia pneumoniae*. Starting with sera from infected patients, blotting against clinical isolates of *Chlamydia pneumoniae* showed the presence of an immunodominant antigen with an apparent molecular weight of 51 kDa, the antigen being stable to and released by octylglucoside treatment. N-terminal amino acid sequencing of the protein of the 51 kDa band allowed sequence database probing, in turn identifying a *C. pneumoniae* protein and a *C. trachomatis* homologue. Epitope mapping allowed the identification of antigenic peptides, which together with antibody against them were tested for their therapeutic and diagnostic efficacy.

Western Blotting - Using the Novex nuPAGE Electrophoresis System.

1. SDS PAGE

Preparation of Sample:

- 1. 100 μ l of Novex SDS Sample loading buffer was added to 400 μ l of a preparation of a *Chlamydia pneumoniae* clinical isolate and the mixture placed into a boiling waterbath for 10 minutes.
- 2. $10 \mu l$ of the mixture was loaded into each well of a Novex 4-12% Bis-Tris NuPage gel (1.0 mm, 12 well). In addition, 4 μl of Novex Multimark molecular weight standards were added to a single well on each gel.
- 3. Electrophoresis was performed using 1x Novex MOPS electrophoresis buffer at 200v for 40 minutes.

Western Transfer Protocol:

- 1. The blotting apparatus and the gel membrane "sandwiches" were assembled according to the protocol described in the Novex instruction booklet provided with the gels.
- 2. Blotting was performed using 1x Novex Transfer buffer containing 20% methanol. Transfer was carried out at 30v (constant) for 1 hour.
- 3. Following transfer, the membranes were removed from the apparatus and left to "Block" overnight in 3% Bovine Serum Albumin (BSA) at 4 °C.

Probing With Patient's Serum:

- 1. The membranes were cut into strips and placed into the wells of incubation trays. Patients' serum was diluted 1 in 20 in 3% BSA and 2 ml added to each strip. (2 strips per patient).
- 2. The membranes were incubated at room temperature for 2 hours with agitation.
- 3. The strips were washed 5 times over 30 minutes with 0.85% NaCl/0.01% Tween 20.
- 4. 2 ml of goat anti-human IgM or IgG alkaline phosphatase conjugated antiimmunoglobulin diluted 1 in 4000 in 3% BSA were added to each strip. The strips were incubated for a further hour at room temperature with agitation.
- 5. The membranes were washed a further 5 times as previously described.

- 6. Antibody-antigen interaction was visualised by the addition of NBT/BCIP (50 mg/ml) in pH 9.5 phosphate buffer.
- 7. The reaction was allowed to proceed until the bands had reached the required intensity.

Sera

- Group A: Children with respiratory tract infection and no evidence of *Chlamydia* pneumoniae as shown by negative microimmunofluorescence (less than 1 in 64) test (n=19).
- Group B: Children with respiratory tract infection and a microimmunofluorescence titre greater than 1 in 512 (n=18).
- Group C: Patients undergoing cardiac surgery for advanced coronary disease (n=32).

 Ten of these had antibody on immunoblot.
- Group D: Adults with respiratory tract infection and a chlamydia complement fixation test greater than 1 in 40 (n=27) using LGV 2 as an antigen.
- Group E: Adults with pelvic inflammatory disease due to *Chlamydia trachomatis* (n=21).
- Group F: Sera (n=11) which were positive for the 60/62 kDa doublet and band at 51 kDa were retested on antigen prepared from *Chlamydia pneumoniae* where the purified elementary bodies were incubated with 1% octylglucoside at 37 °C for 30 minutes rather than in SDS.

WO 00/46359 PCT/GB00/00237

- 11 -

Results:

Results of the sera blotting experiments are shown in Table 1. It should be noted that sera blotting determines the presence in patients of antibodies specific against a given antigen, and so when a patient has previously been infected by a pathogen and developed an immune response against an antigen, that immune response may still be detectable at a later date when the patient is no longer infected. Hence background results must be interpreted in light of the general infection of a population by the pathogen. For example, the general population has an infection rate by adulthood of approximately 10% for *C. pneumoniae*, thus a background rate of detection of *C. pneumoniae* antigens of up to 10% should be expected.

Conclusions:

The sera from Group A children did not recognise *C.pneumoniae* on immunoblot. The Group B sera from children with evidence of *C.pneumoniae* infection recognised a range of antigens with apparent molecular weights ranging from 30 to 180 kDa. IgM for an antigen complex at 60/62 kDa which occurred as a doublet was immunodominant as well as an antigen at 51 kDa. For IgG the antibody was most pronounced for the antigen at 51 kDa. In the cardiac patients, 23 produced antibody and this was for IgM against the bands at 67, 60/62 and 51 kDa. For IgG this was the band at 51 kDa. For Group D IgM was most pronounced for the 60/62 kDa doublet and IgG for the band at 180 kDa and the doublet at 60/62 kDa. This group of sera contains those with infection most likely due to *Chlamydia psittaci*. The sera from Group E patients infected with *Chlamydia trachomatis* did not cross-react.

Group F Sera

On re-blotting with those sera previously positive for the 60/62 kDa doublet and 51 kDa, the doublet disappeared whilst the band at 51 kDa remained. This showed that the band at 51 kDa was stable to and released by octylglucoside treatment.

WO 00/46359 PCT/GB00/00237

- 12 -

Solubility in Octylglucoside

Using samples from Group F patients, separation of antigens from elementary bodies using 1-D gel electrophoresis and SDS gave a different staining pattern compared to using 1-D gel electrophoresis and octylglucoside. The 51 kDa band was still visible after octylglucoside. The pair of antigenic bands at 60/62 kDa was not visible in octylglucoside. Therefore a distinguishing character of the 51 kDa antigen of the present invention is its solubility in octylglucoside.

N-Terminal Amino Acid Sequencing

N-Terminal amino-acid sequencing was performed upon the 51 kDa band. The resulting sequence was then used to query the Chlamydia Genome Project database which identified the protein of SEQ ID NO: 2 and a *C. trachomatis* homologue.

Epitope Mapping

A series of overlapping peptides of 15 amino acids covering the derived amino acid sequence of the protein were synthesised on polyethylene pins with reagents from an epitope scanning kit (Cambridge Research Biochemicals, Cambridge, UK) as described previously by Geysen *et al.* (1987, Journal of Immunological Methods, 102: 259-274). Peptide 1 consisted of residues 1 to 15, peptide 2 consisted of residues 2 to 16 etc. The reactivity of each peptide with patient sera (diluted 1:200) was determined for IgG by ELISA. Data were expressed as A405 after 30 minutes of incubation.

Sera from patients as follows:

- Group 1: Children with respiratory tract infection and no evidence of *Chlamydia* pneumoniae as shown by negative immunoblot and microimmunofluorescence (less than 1 in 64) (n = 3).
- Group 2: Children with respiratory tract infection, positive immunoblot and microimmunofluorescence test greater than 1 in 512 (n = 6).
- Group 3: Patients undergoing cardiac surgery for advanced coronary disease and antibody on immunoblot (n = 2).
- Group 4: Patients presenting with history of chest pain, negative troponin (<0.2), negative immunoblot (n = 3).
- Group 5: Patients presenting with early coronary, positive troponin (>0.2) and antibody on immunoblot (n = 8).

Results

Epitope mapping

Epitope mapping defined eleven areas where children with acute chlamydial infection produced wells with a mean optical density (OD) greater than 1. In the case of epitopes having SEQ ID NOs: 4, 5, 6, 7, 8, 10, 12 and 14 the mean OD was at least 2 standard deviations above that of Group 1 (children with no evidence of *C.pneumoniae* infections). This applied also to Groups 3, 4 and 5 with the exception of SEQ ID NO: 5 which was positive in Groups 4 and 5.

Peptide 1(SEQ ID NO: 15) representing epitope having the sequence of (i.e. which is carried by the peptides having the sequence of) SEQ ID NO: 8 and peptide 2 (SEQ ID NO: 16) representing the carboxy end of SEQ ID NO: 4, the epitope having the sequence of SEQ ID NO: 5 and the amino end of SEQ ID NO: 6 were synthesised.

Preparation of rabbit polyclonal serum

New Zealand white rabbits were pre-bled and then immunised subcutaneously with either peptide 1 or peptide 2 (0.1 ml of 1 mg/ml) conjugated to KLH suspended in either Freund's adjuvant (injection at day 0) or Freund's incomplete adjuvant on days 14, 42, and 70). Serum was obtained for indirect ELISA at the terminal bleed-out.

Indirect ELISA

By a simple adsorption of each peptide to a microtitre plate the following procedure was performed The peptide was dissolved in 2 ml of 0.01 M phosphate buffer saline (PBS), pH 7.2 and diluted to a concentration of 10 μ g/ml (1/100) in the same buffer.

- 150 μl aliquots of peptide (10 μg/ml in 0.01 M PBS) were pipetted into the wells of a Falcon 3912 microassay plate and were incubated overnight at 4 °C.
- 2. The unbound peptide was removed by washing four times (4 x 10 minutes) with 0.05% Tween 20 in 0.01 M PBS (pH 7.2).
- 3. The plates were blocked with 2% skimmed milk-10% FCS in 0.01 M PBS for 1 hour at 37 °C.
- 4. The plates were washed four times (4 x 10 minutes) with 0.05% Tween 20 in 0.01 M PBS and the serum under investigation was added (1/100 dilution in blocking solution) into the wells of micro assay plate (three wells used for each serum) and incubated for 2 hours at 37 °C.
- 5. The plates were washed four times (4 x 10 minutes) with 0.05% Tween 20 in 0.01 M PBS and secondary antibody, anti-rabbit IgG peroxidase conjugate (1/1000 dilution in blocking solution) was added and incubation proceeded for 1 hour at 37 °C.
- 6. The plates were washed four times (4 x 10 minutes) with 0.05% Tween 20 in 0.01 M PBS, followed by a further washing with 0.01 M PBS. The plate was then incubated for 45 minutes at room temperature with agitation in 0.5

- mg/ml of freshly prepared 2,2 Azino-bis [3-ethylbenz-thiazoline-6-sulfonic acid] diammonium (ABTS tablets) in pH 4.0 citrate buffer with 0.01% (w/v) hydrogen peroxide.
- 7. Optical density (OD) measurements were made with an ELISA plate reader (Titertek Miltiscan) at a wavelength of 405 nm.
- 8. The average readings for each three wells for each serum was determined.

Results

The results shown in Table 3 demonstrate seroconversion to each individual peptide.

Expression of the amino-end of the protein

The sequence was codon optimised (Genosys, California) for *E.coli* and a BamHI and Not1 site added to opposite ends. The optimised sequence and PET 29 vector (Novagen, Wisconsin) were restriction digested using BamHI and Not1 and transformed by heat shock into *E.coli* strain BL21 (Invitrogen, Carlsbad, California). The expressed amino acids were from amino acids 1-292 and included the epitopes represented by peptides 1 and 2. This construct included an S-tag and Thrombin cleavage site at the amino end and histidine tag at the carboxy end (SEQ ID NO: 3).

Purification

The transformants were expressed as follows. Briefly, 5 ml of an overnight culture was used to inoculate 500 ml LB (50 μg/ml kanamycin, 34 μg/ml chloramphenicol) which was grown for 2 hours at 37 °C to an OD 600 of 0.5, then induced for 3 hours with 0.1 mM IPTG (Sigma, Poole Dorset). The cells were pelleted and disrupted by crushing at -20 °C in an XPRESS. The buffer (50 mm NaH₂PO₄, 0.5 M NaCl, 10 mm imidazole) and the cell debris pelleted down. The supernatant was filter sterilised and put on a Ni-NTA agarose slurry affinity column (Qiagen) in order to capture the His-tagged recombinant protein. The column was washed 3 times with 4 ml of washing buffer and

WO 00/46359 PCT/GB00/00237

- 16 -

the protein eluted maximally with 150 mM imidazole. The protein gave a single band on a 10% acrylamide gel stained with Coomassie Brilliant Blue with an apparent molecular weight of 37 kDa. On Western blot counterstaining with the anti-His mouse alkaline phosphate conjugate (1:2,500) (Sigma, Dorset, Poole) this produced a single band at 37 kDa and a breakdown product at 35 kDa. The protein concentration of the elute was measured and standardised to 10 mg/ml.

Amino acid sequencing

The protein was amino end cleared to remove the S-tag using a Thrombin cleavage Kit (Novagen). The digestion reaction was 5 μ l 10 x Thrombin cleavage buffer, 0.5 mg purified recombinant protein, 1 μ l of 0.01 μ g/ml Thrombin which was left at room temperature for 18 hours. The reaction mix was run on a 12% SDS-PAGE gel and transferred onto PVDF membrane (Amersham, Chalfont, UK). This was stained with Coomassie Brilliant Blue and the protein bands destained and excised. Direct amino acid sequencing gave amino acids 28-32 of SEQ ID NO: 3 which matched the amino end (Department of Biochemistry, University of Cambridge).

Human recombinant antibodies

These peptides and the purified recombinant proteins were used to pan the phage display library. The peptide and recombinant protein were used at 10 mg/ml on NunC immunotubes Bst-N1 fingerprints of the PCR-amplified ScFv inserts before panning showed a highly heterogeneous library. After panning against peptide 1, 7 fingerprints were identified of which four were represented by more than one clone (A, B, C, D). These were combined as a pool for a neutralisation assay (pool 1) (below). After panning against peptide 2, clone A was present as well as a new ScFv, E. A and E were combined to produce pool 2. Against the clone recombinant fragment ScFvs E, F and G were present as well as a further ScFv, H. ScFvs E, F, G and H were tested together as pool 3.

Neutralisation assays

Chang cells (50 ml of 10^6 cells/ml) in maintenance media were grown overnight at 37 °C with 5% CO₂. Chang cells (1 ml of $1x10^6$ cells/ml maintenance media) were grown overnight at 37 °C with 5% CO₂ in plastic bijoux containing a thin glass circle on which the cells can grow. For recombinant protein or peptide assay (0.1 μ l/ml), 100 μ l of each sample was incubated with shaking for 1 hour with the cells at 37 °C. For the phage and sera assays, 100 μ l of each sample (1:10 rabbit sera or dialysed phage pools 1-3) were incubated with 100 μ l elementary bodies (EB) for 1 hour at 37 °C, shaking. After this first incubation, the 100 μ l EB or 200 ml of the phage or rabbit sera/EB mix was added to the Chang cells. This was incubated with shaking for 1 hour at 37 °C. The supernatant was removed from every sample and replaced by 1 ml of fresh maintenance media. This was incubated at 37 °C with 5% CO₂ for 72 hours.

For both assays, the inclusion bodies were fixed and stained the following way; the cells were washed twice with PBS, then fixed with 100% methylated spirits for 10 minutes and washed twice again with PBS. The glass circles were incubated for 30 minutes with 10 µl of mouse *C.pneumoniae* inclusion bodies monoclonals (Mab) then washed 3 times with PBS and incubated for 30 minutes with 100 µl of fluorescein conjugated anti-mouse IgG. The inclusion bodies were then observed by fluorescence microscopy and three 200X fields counted. EB only samples were used as a positive control for chlamydial infection and dialysed phage supernatant without EB as a negative control.

Results

See Table 4 (Table of Neutralisation Assays).

Conclusion

Pre-incubation with the rabbit antiserum against peptide 2 and peptide 2 itself reduced the infectivity due to *C.pneumoniae*. Incubation with peptide 1 produced a similar reduction. The pools of phages were also active.

Overall this demonstrated the immunogenicity of the antigen, the potential therapeutic effect of peptides representing its key epitopes and both rabbit hyperimmune antiserum and ScFvs against these epitopes.

Table 1

Apparent	Gro	oup B	Gro	up C	Gro	ın D	Gro	up E
Molecular	İ	=18)		=18)	(N=			=21)
Weight (kDa)	IgM	IgG	IgM	IgG	IgM	IgG	IgM	IgG
180	1	2		2	1	6		1
130		2			1	4		
120	1	5		1	1	5		1
98		5		1	2	5		2
90		2				2		
67		2	5	1			1	1
60/62*	8	5	5		13	7	2	2
51	7	11	9	10	2	3	1	2
47	1	1	1		0	0	0	0
40	0	0	0	3	0	0	0	1
30		4	0	3		2		2

^{*} runs as a doublet within 1 mm of each other

Table 2

		- Value for ^a								
Well	Epitope	Group 1	Group 1 Group 2 Group 3 Group 4		Group 5					
No.	SEQ ID NO	(n = 3)	(n = 6)	(n = 2)	(n = 3)	(n = 8)				
3	9	0.538±0.205	1.028±0.423	0.425±0.036	0.416±0.184	0.499±0.191				
4		0.599±0.252	1.487±0.462	0.502±0.036	0.407±0.107	0.438±0.162				
13	10	0.462±0.203	1.103±0.229	0.473±0.026	0.421±0.162	0.427±0.188				
31	11	0.491±0.192	1.103±0.310	0.440±0.004	0.407±0.105	0.310±0.129				
41	12	0.547±0.235	1.169±0.256	0.474±0.024	0.393±0.08	0.376±0.158				
43	13	0.598±0.258	1.223±0.323	0.558±0.015	0.423±0.119	0.406±0.181				
55	4	0.547±0.235	1.265±0.334	0.475±0.02	0.373±0.076	0.381±0.042				
58	5	0.611±0.019	1.025±0.06	0.611±0.019	1.127±0.253	0.800±1.232				
59	6	0.494±0.166	1.096±0.267	0.547±0.009	0.546±0.200	0.702±0.144				
60	7	0.489±0.129	1.048±0.270	0.483±0.064	0.388±0.008	0.449±0.140				
61		0.530±0.236	1.051±0.262	0.59±0.089	0.446±0.09	0.784±0.257				
76	8	0.485±0.158	1.174±0.255	0.654±0.068	0.564±0.223	0.666±0.266				
79	14	0.510±0.235	1.21±0.273	0.418±0.003	0.423±0.127	0.388±0.153				

^a Optical density ± Standard deviation

 a Pre Serum
 Post Serum

 Peptide 1
 0.055 ± 0.01 0.591 ± 0.06

 Peptide 2
 0.056 ± 0.01 0.507 ± 0.04

- 21 -

<u>Table 4</u> - Table of Neutralisation Assays

·	Number of Elementary Bodies in Three				
	200x Fields				
Cell control (dialysed phage	0				
supernatant)					
Cell control (elementary bodies)	30				
Rabbit anti-serum					
Versus peptide 1	30				
Versus peptide 2	19				
Pre-incubation					
Peptide 1	13				
Peptide 2	0				
Recombinant protein	12				
Phage Pools					
Pool 1	18				
Pool 2	N/D				
Pool 3	21				

^a optical density ± standard derivation

CLAIMS

- 1. A *C.pneumoniae* protein having the amino acid sequence of SEQ ID NO: 2 for use in a method of treatment or diagnosis of the human or animal body.
- 2. A nucleotide sequence encoding a protein according to claim 1 for use in a method of treatment of the human or animal body.
- 3. A nucleotide sequence according to claim 2, having the sequence of SEQ ID NO: 1.
- 4. The use of a protein, immunogenic fragment thereof or nucleotide sequence encoding same according to any one of the preceding claims in the manufacture of a medicament for the treatment of infection due to *C.pneumoniae*.
- 5. The use of an immunogenic fragment according to claim 4, having the amino acid sequence of any one of SEQ ID NOs: 4-14 in the manufacture of a medicament for the treatment of infection due to *C.pneumoniae*.
- 6. The use of an inhibitor specific against the protein, immunogenic fragment or nucleotide sequence encoding same according to any one of the preceding claims in a method of manufacture of a medicament for the treatment of infection due to *C.pneumoniae*.
- 7. The use of an inhibitor according to claim 6, the inhibitor being selected from the group of an antibody, DNA vaccine, ribozyme and antisense oligonucleotide.

- 8. A method of manufacture of a medicament for the treatment of infection by *C.pneumoniae* characterised in the use of a protein, immunogenic fragment thereof or nucleotide sequence encoding same according to either one of claims 4 or 5.
- 9. A method of manufacture of a medicament for the treatment of infection due to *C.pneumoniae* characterised in the use of an inhibitor according to either one of claims 6 or 7.
- 10. The use of a protein according to claim 1 or an immunogenic fragment thereof or a binding agent specific to same or an inhibitor of same in the manufacture of a diagnostic test for *C.pneumoniae*.
- 11. A kit of parts for a diagnostic test for *C.pneumoniae*, characterised in that it comprises a protein according to claim 1 or an immunogenic fragment thereof or a binding agent specific to same or an inhibitor of same.
- 12. A diagnostic test method for infection due to *C.pneumoniae* comprising the steps of:
 - reacting an antibody specific against the protein according to claim
 with serum from a patient;
 - ii) detecting an antibody antigen binding reaction; and
 - iii) correlating the detection of an antibody antigen binding reaction with the presence of the protein.

- 13. A diagnostic test method according to claim 12, being a method of diagnosis of the human or animal body.
- 14. A method of treatment of infection due to *C.pneumoniae* comprising the step of administering to a patient a medicament comprising a protein, immunogenic fragment thereof, nucleotide sequence encoding same or an inhibitor thereof according to any one of claims 4-7.

FOR UTILITY/DESIGN OIP/PCT NATIONAL/PLANT ORIGINAL/SUBSTITUTE/SUPPLEMENTAL DECLARATIONS

RULE 63 (37 C.F.I \$33) DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

PW FORM

DECLARATIONS IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

As a below named inventor, I hereby declare that my residence, post office address and citizenship are a stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a natent is sought on the INVENTION ENTITY OF New York or the INVENTION ENTITY OF NEW YORK OF THE INVENTION ENTITY
	ot matter willen				jiit on the <u>ir</u>	NVENTION EN	IIILED Me	edicamer	nt	
the sp X A. □	ecification of w is attached here	hich (CHECK	applicable Bo	OX(ES))						
BOX(ES) → E	3. 🔲 was filed o	on		as	U.S. Appli	cation No	1			
\rightarrow \rightarrow \bigcirc	C. 🔲 was filed a	as PCT Inte	rnational A	pplication N	lo. PCT/	GB00/00237	on	28 Jar	nuary 2000	
and (if applicable to I hereby state that I ha	ive reviewed and	understand the	contents of the	above identified	specification	including the elei-	ne ac ame:			
foreign priority benefits Application which desi	s under 35 U.S C. gnated at least or ernational Applicat	119(a)-(d) or 3 ne other country	65(b) of any fore than the United	eign application(s I States, listed be	patentability b) for patent or elow and have	as defined in 37 (r inventor's certific e also identified be	F.R. 1 56. ate, or 365(Except as a) of any Po		claım
PRIOR FOREIGN	APPLICATION	(<u>S)</u>			Date fire	st Laid-	Date Pa	tented		
<u>Number</u> 9902555.3	<u>Country</u> GB		y/MONTH/Ye /02/1999	ar Filed		or Published		ranted	Priority NOT Cl	aimed
application is in addition defined in 37 C.F.R. 1. application: PRIOR U.S. PROV Application No. (s	r, I hereby claim dications listed about the thick that discloses 56 which became ISIONAL, NON eries code/ser	omestic priority we or below an d in such prior a available betw PROVISION. ial no.)	benefit under 3 d, if this is a con applications, I ac een the filing da AL AND/OR F Day/MON	5 U.S C. 119(e) trinuation-in-part the kinowledge the content of the of each such part of the content of the co	or 120 and/or (CIP) application of the disclosorior application of the theorem (CIP) and	tion, insofar as the e all information kon and the national pending, all	e subject manown to me all or PCT int	tter disclos to be mate ernational f	filing date of this Priority NOT Cla	<u>timed</u>
And I hereby appoint Persons of that firm whereby appoint Persons of that firm whereby appoint Persons no letter person/assignee/at Tisclosure to be represused USE ONL	rillsbury Winthrop o are associated of the Patent and Tonger with their fir torney/firm/ organ ented unless/until	LLP, Intellectua with USPTO Cu rademark Office m, to add new p	al Property Groupstomer No 909 econnected their persons of their lich first sends/sebove Firm and/o	p, telephone nun (see below labe rewith and with the Firm to that Cust	nber (703) 90 d) individually ne resulting pa omer No , and	5-2000 (to whom a and collectively m atent, and I hereb d to act and rely o	all communic ny attorneys y authorize t n instruction	cations are to prosecut hem to dele	to be directed), and te this application and ete from that Custome	io · No with
1) INVENTOR'S SI	GNATURE:	fa	n P-1	h			1,10	L i.	200 1	
	ames	(1)	P	#	1 51151	Date:	"/ 	74.Y	001	
1.29		First	I P	Middle Initial	BURNI		\ 	<u>, , , , , , , , , , , , , , , , , , , </u>		
Residence A	derley Edge		BL	Middle Initial GB				y Name		
	dono, Lugo	City			tota/Caraina C		GB			
Mailing Address	1 Grevstoke		ev Edge, Che	shire, SK9 7F	ate/Foreign C	ountry		Cou	untry of Citizenship	
include Zip Code)	· Groyotoke	Dive, Alden	ey Lage, One	Silite, SN9 7P	Y, GB					
тр оссо)				, -						
2) INVENTOR'S SI	GNATURE:	MIN	lattle	<u> </u>		Date:	11 50	Lhy ?	2001	
lame Ri	uth		<u>. c</u>	- mpy - TONE diktor	MATTH	EWS		'		
		First		Middle Initial			Family	/ Name		
Residence Al	derley Edge	G	BL	GB			GB			
		City			ate/Foreign Co	ountry		Cou	intry of Citizenship	
/lailing Address	1 Greystoke	Drive, Alderle	ey Edge, Che	shire, SK9 7P					, c. c.meonomp	
include Zip Code)						· · · · · · · · · · · · · · · · · · ·				
FOR ADDITION See addition	ONAL INVEI	NTORS se orities on a	e attached ttached pag	page. ge (incorpor	ated here	in by referer Atty. Dkt.	nce). No. P	(Ma	#)	

Rule 56(a) & (b) = 37 C.F.R. 1.56(a) & (b) PATENT AND TRADEMARK CASES - RULES OF PRACTICE DUTY OF DISCLOSURE

(a) ... Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the [Patent and Trademark] Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability...(b) information is material to patentability when it is not cumulative and (1) It also establishes by itself, or in combination with other information, a prima facie case of unpatentability of a claim or (2) refutes, or is inconsistent with, a position the applicant takes in: (i) Opposing an argument of unpatentability relied on by the Office, or (ii) Asserting an argument of patentability

PATENT LAWS 35 U.S.C.

§102. Conditions for patentability; novelty and loss of right to patent

A person shall be entitled to a patent unless--

- (a) the invention was known or used by others in this country, or patented or described in a printed publication in this or a foreign country, before the invention thereof by the applicant for patent or
- (b) the invention was patented or described in a printed publication in this or a foreign country or in public use or on sale in this country, more than one year prior to the date of the application for patent in the United States, or
- (c) he has abandoned the invention, or
- the invention was first patented or caused to be patented, or was the subject of an inventor's certificate, by the applicant or his legal representatives or assigns in a foreign country prior to the date of the application for patent in this country on an application for patent or inventor's certificate filed more than twelve months* before the filing of the application in the United States, or
 - the invention was described in a patent granted on an application for patent by another filed in the United States before the invention thereof by the applicant for patent, or on an international application by another who has fulfilled the requirements of paragraphs (1), (2), and (4) of section 371(c) of this title before the invention thereof by the applicant for patent, or
 - he did not himself invent the subject matter sought to be patented, or
 - before the applicant's invention thereof the invention was made in this country by another who had not abandoned, suppressed, or concealed it. In determining priority of invention there shall be considered not only the respective dates of conception and reduction to practice of the invention, but also the reasonable diligence of one who was first to conceive and last to reduce to practice, from a time prior to conception by the other.

§103. Condition for patentability; non-obvious subject matter

- (a) A patent may not be obtained though the invention is not identically disclosed or described as set forth in section 102 of this title, if the differences between the subject matter sought to be patented and the prior art are such that the subject matter as a whole would have been obvious at the time the invention was made to a person having ordinary skill in the art to which said subject matter pertains. Patentability shall not be negatived by the manner in which the invention was made. . . .
- (c) Subject matter developed by another person, which qualified as prior art only under subsection (f) or (g) of section 102 of this title, shall not preclude patentability under this section where the subject matter and the claimed invention were, at the time the invention was made, owned by the same person or subject to an obligation of assignment to the same person.

(f)

T.

^{*} Six months for Design Applications (35 U.S.C. 172).

United States Patent & Trademark Office

Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

Page(s) of Chemswittel. were not present for scanning. (Document title)

Page(s) of were not present for scanning. (Document title)

□ Scanned copy is best available.